CLAIMS

We claim:

1. An array composition comprising:

5   a) a substrate with a surface comprising discrete sites; and

b) a population of microspheres comprising at least a first and a second
subpopulation, wherein each subpopulation comprises:

i) a bioactive agent; and

ii) an identifier binding ligand that will bind a decoder binding ligand such

10   that the identification of the bioactive agent can be elucidated;

wherein said microspheres are distributed on said surface.

2. An array composition comprising:

a) a substrate with a surface comprising discrete sites; and

15   b) a population of microspheres comprising at least a first and a second
subpopulation, wherein each subpopulation comprises a bioactive agent and does not
comprise an optical signature, wherein said microspheres are distributed on said
surface.

3. A composition according to claim 1 or 2 further comprising at least one decoder binding
20   ligand.

4. A composition according to claim 1 or 2 wherein said bioactive agents are nucleic acids.

25   5. A composition according to claim 1 or 2 wherein said bioactive agents are proteins.

6. A method of making a composition comprising:

a) forming a surface comprising individual sites on a substrate;

b) distributing microspheres on said surface such that said individual sites contain

30   microspheres, wherein said microspheres comprise at least a first and a second
subpopulation each comprising a bioactive agent and do not comprise an optical
signature.

41

7. A method of making a composition comprising:

a) forming a surface comprising individual sites on a substrate;

b) distributing microspheres on said surface such that said individual sites contain microspheres, wherein said microspheres comprise at least a first and a second subpopulations each comprising:

i) a bioactive agent; and

ii) an identifier binding ligand that will bind a decoder binding ligand such that the identification of the bioactive agent can be elucidated.

8. A method of decoding an array composition comprising

a) providing an array composition comprising:

i) a substrate with a surface comprising discrete sites; and

ii) a population of microspheres comprising at least a first and a second subpopulation, wherein each subpopulation comprises a bioactive agent;

wherein said microspheres are distributed on said surface;

b) adding a plurality of decoding binding ligands to said array composition to identify the location of at least a plurality of the bioactive agents.

9. A method according to claim 8 wherein at least one subpopulation of microspheres comprises an identifier binding ligand to which a decoding binding ligand can bind.

10. A method according to claim 8 wherein said decoding binding ligands bind to said bioactive agents.

11. A method according to claim 8 wherein said decoding binding ligands are labeled.

12. A method according to claim 8 wherein the location of each subpopulation is determined.

13. A method of determining the presence of a target analyte in a sample comprising:

a) contacting said sample with a composition comprising:

i) a substrate with a surface comprising discrete sites; and

ii) a population of microspheres comprising at least a first and a second

subpopulation each comprising a bioactive agent and do not comprise an optical signature;

   wherein said microspheres are distributed on said surface such that said discrete sites contain microspheres; and

5   b) determining the presence or absence of said target analyte.

14. A method of determining the presence of a target analyte in a sample comprising:

   a) contacting said sample with a composition comprising:

      i) a substrate with a surface comprising discrete sites; and

10   ii) a population of microspheres comprising at least a first and a second subpopulation each comprising:

         1) a bioactive agent; and

         2) an identifier binding ligand that will bind a decoder binding ligand such that the identification of the bioactive agent can be elucidated;

15   wherein said microspheres are distributed on said surface such that said discrete sites contain microspheres; and

   b) determining the presence or absence of said target analyte.

add
b4
ADD
E∂